



UNIVERSITI PUTRA MALAYSIA

**UTILISATION OF FOOD WASTES FOR THE PRODUCTION OF
ORGANIC ACIDS AND BIOPLASTICS**

NORRIZAN ABDUL WAHAB

FSMB 2003 22

**UTILISATION OF FOOD WASTES FOR THE PRODUCTION OF
ORGANIC ACIDS AND BIOPLASTICS**

By

NORRIZAN ABDUL WAHAB

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of Requirements for the Degree of Master of Science**

June 2003



ALLAH
MY DESTINY IS HERE
DROP DOWN AND STRONG AGAIN
IT SO MEANINGFUL SPIRIT TO
UNDER YOUR BLESS
I'M SUCCESSFUL
ALHAMDULILLAH... THANK TO
FOR ALL THINGS
ITS

MY BELOVED PARENTS,
ABDUL WAHAB & ZAIMAH
THANK YOU FOR EVERYTHING
ANYTHING YOU DO
YOUR ENCOURAGE, YOUR GIVING, YOUR WISHING AND ALL
ALHAMDULILLAH... I LOVE YOU SO MUCH

ABANG, KAK, NYAH, YANG, AND YONG,
FOR THERE WISHING ME ALL THE BEST...
ALHAMDULILLAH... SO APPRECIATED

ADIK
AL-FATIHAH (28/7/2001)
DEEPLY MISSED BY FAMILY...
ALWAYS LOVED AND REMEMBERED...
MAY SHE REST IN PEACE
AMIN...

MY BELOVED,
HAMBALI
HAPPY AND SAD WITH ME
YOUR WISHING WITH ME
TO ENCOURAGE ME
TRUST AND LOYALTY
HAS MADE ME FEEL VERY LUCKY
ALHAMDULILLAH... INVALUABLE LOVE

FRIENDS
YOUR HELPFUL AND YOUR ENCOURAGE
IT SO MEANINGFUL TO ME
I'M VERY APPRECIATE
THANK YOU
FOR ALL..

Abstract of a thesis presented to the Senate of Universiti Putra Malaysia in Fulfillment of the requirement for the degree of Master of Science.

UTILISATION OF FOOD WASTES FOR THE PRODUCTION OF ORGANIC ACIDS AND BIOPLASTICS

By

NORRIZAN ABDUL WAHAB

October 2003

Chairman: Professor Dr. Mohd Ali Hassan

Faculty: Food Science and Biotechnology

In this study, food wastes were used as substrates for the production of polyhydroxyalkanoate (PHA) by *Ralstonia eutropha* (formerly known as *Alcaligenes eutrophus*). Restaurant waste obtained from a local restaurant and house was blended with an equal weight of water. PHA production was achieved via a two-stage process; organic acids production from food waste followed by PHA production from the organic acids. In the first stage, the study focused on the production of short-chain organic acids from complex organic fraction of food waste. The food waste was subjected to anaerobic treatment by the natural micro flora present in a 2-liter batch stirred tank bioreactor with mixing at 200 – 400 rpm. The effect of pH, temperature, substrate compositions of food waste on the production of volatile fatty acids (VFA) was evaluated. For the effect of different pH, the batch anaerobic medium was done [uncontrolled pH, controlled at pH 7 for different number of days (initial, one day, two days and whole duration)].

The effect of different temperatures (30°C, 37°C, 45°C) and different substrate compositions [kitchen waste (A), restaurant waste (B) and restaurant waste mixed with fruits waste (C)] have been done in the research. Restaurant waste mixed with fruit waste (substrate C) produced 58 g/L organic acids compared to 42 g/L from kitchen waste (substrate A) under same conditions at 37°C without control pH. Lactic acid was the most predominant acid, followed by acetic, propionic, butyric and formic acids. The study on the effect of temperature with uncontrolled pH shows that 37°C is the best temperature, producing 58 g/L organic acids, with 83% lactic acid. In the effect of pH, it was found that the highest amount of organic acids produced was 68 g/L, with 84% lactic acid when the initial pH was adjusted to 7.0 without further pH control.

Overall the best conditions for organic acids production is pH controlled at 7 in the initial treatment at 37°C using restaurant wastes containing fruit waste. Studies on microbial and enzyme activity was done on treated food waste to determine the factors contributing to the high organic acids especially lactic acid. From the microbes identified, *Lactobacillus* was predominantly involved in the fermentation to produce lactic acid. The enzyme lipase, protease, cellulase and α -amylase were detected throughout the treatments leading to the conversion of organic complex to sugar and subsequently to organic acids with α -amylase as the highest activity. In the recovery of organic acids from the fermented food waste, centrifugation was carried out to remove the solids and oils. The supernatant was collected and concentrated by evaporation for the second stage of bioplastic production.

The results show that by evaporation method, the organic acids were concentrated two-fold to about 100 g/L for use as substrates for fed-batch fermentation. PHA production from organic acids fermented from food waste, as a carbon source for fed-batch fermentation is 4.2 g/L with 88% of dry cell weight. From the experiment, C/N 10-20 because ammonium concentration in the original food waste was high. This encouraged cell growth in the medium rather than PHA production. In the PHA production, since C/N must be high (around 100-150), removal of ammonium was successfully achieved using Dowex 88wx resin for improved PHA production. The ammonia-deficient organic acids from food wastes were successfully converted to PHA by *Ralstonia eutropha* strain ATCC 17699, producing 8.9 g/L with 90%(g/g) PHA content.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

PENGGUNAAN SISA MAKANAN BAGI MENGHASILKAN ASID ORGANIK DAN BIOPLASTIK

Oleh

NORRIZAN ABDUL WAHAB

Jun 2003

Pengerusi: Professor Mohd Ali Hassan, Ph.D.

Fakulti: Sains Makanan dan Bioteknologi

Dalam kajian ini, sisa makanan digunakan sebagai substrat bagi penghasilan polyhydroxyalkanoate (PHA) oleh *Ralstonia eutropha* (juga dikenali sebagai *Alcaligenes eutrophus*). Sisa makanan yang diperolehi daripada restoran tempatan dan rumah kemudian ditimbang dengan jumlah berat air yang sama untuk dihancurkan (blender). Terdapat dua peringkat kajian bagi penghasilan PHA iaitu penghasilan asid organik dari sisa makanan dan diikuti dengan penghasilan PHA menggunakan asid organik yang diperolehi sebagai substrat. Dalam tahap pertama, kajian difokuskan kepada penghasilan rantaian pendek asid organik daripada pemecahan organik kompleks dari sisa makanan tersebut. Komposisi sisa makanan yang hampir sama bagi rawatan anaerobic dipengaruhi oleh kehadiran microflora semulajadi di dalam 2-liter bioreactor tangki berpengaduk yang dicampur pada kelajuan dari 200 hingga 400 rpm.

Kajian ke atas perbezaan pH [tanpa kawalan pH, pH dikawal pada pH 7 (awal rawatan, satu hari, dua hari dan sepanjang rawatan)], perbezaan suhu (30°C, 37°C, 45°C) dan komposisi substrat yang berbeza [sisa dapur (A), sisa restoran (B) dan sisa restoran yang mengandungi sisa buah (C)] dari sisa makanan dalam memberikan kesan ke atas penghasilan asid lemak bebas di nilai. Substrat C dari sisa restoran yang mengandungi sisa buah menghasilkan 58 g/L berbanding dengan substrate A dari sisa rumah yang hanya menghasilkan 42 g/L asid organik dalam keadaan yang sama iaitu pada suhu 37°C tanpa pengawalan pH. Umumnya, asid laktik merupakan asid yang paling banyak dihasilkan diikuti oleh asetik, propionik, butirik dan asid formik. Dalam kajian ke atas kesan suhu, suhu pada 30°C, 37°C dan 45°C digunakan tanpa kawalan pH. Didapati keputusan menunjukkan suhu 37°C merupakan suhu yang paling optimum menghasilkan 58 g/L asid organik dengan 83% asid laktik.

Bagi kesan pH, media di biarkan tanpa kawalan pH dan pengawalan pada pH 7 untuk bilangan hari yang berbeza. Didapati penghasilan asid organik yang tinggi iaitu 68 g/L, 84% asid laktik diperolehi, apabila pada awal rawatan, pH di kawal pada pH 7 dan seterusnya dibiarkan tanpa pengawalan pH. Dari keseluruhan keputusan, keadaan yang paling baik untuk penghasilan asid organik ialah dengan mengawal pH pada pH 7 hanya pada awal rawatan pada suhu 37°C dengan menggunakan substrat C. Kajian dalam microbiologi dan aktiviti enzim telah dibuat ke atas fermentasi sisa makanan bagi menentukan faktor yang menyumbang kepada kepekatan asid organik yang tinggi terutama asid laktik.

Didapati microb *Lactobacillus* memainkan peranan penting dalam fermentasi sisa makanan secara anaerobik menghasilkan produk laktik asid. Enzim lipase, protease, cellulase dan α -amylase telah dikesan sepanjang rawatan dalam proses pertukaran kompleks organik kepada gula dan seterusnya ditukarkan kepada asid organik dengan enzim α -amylase mempunyai tahap aktiviti yang tinggi. Bagi memperoleh asid organik daripada fermentasi, sisa makanan diempur bagi membuang pepejal dan minyak. Kemudian supernatan di kumpul dan dipekatkan dengan pemelupaan untuk digunakan bagi penghasilan bioplastik. Hasil dari pemelupaan, asid organik telah dipekatkan dua kali iaitu lebih kurang 100 g/L untuk digunakan sebagai substrat fermentasi suapan sesekelompok.

Penghasilan PHA dari sisa makanan sebagai sumber karbon untuk suapan sesekelompok adalah 4.2 g/L dengan 88% berat sel kering. Didapati nilai C/N adalah rendah (10-20) disebabkan kepekatan amonia di dalam media pengeluaran semakin bertambah semasa fermentasi. Keadaan ini menggalakkan pertumbuhan sel dalam media. Bagi penghasilan PHA, C/N mesti tinggi (100-150), jadi pengurangan kepekatan amonia telah dibuat dengan menggunakan resin Dowex 88wx dalam proses meningkatkan penghasilan PHA. Asid organik dari sisa makanan yang mempunyai kepekatan amonia yang agak rendah telah berjaya ditukarkan kepada PHA oleh *Ralstonia eutropha* ATCC 17699 menghasilkan sebanyak 8.9 g/L dengan 98 % (g/g) kandungan PHA.

ACKNOWLEDGEMENTS

Alhamdulillah, thanks a lot to merciful Allah for giving me the strength to finish this project. Thanks to my supervisors, Prof. Dr Mohd Ali Hassan, Prof Dr. Mohammad Ismail Abdul Karim, Dr Fakhru'l-Razi Ahmadun and Prof. Dr. Yoshihito Shirai, for all their guidance, suggestions and instructions for this project.

My gratitude to all of the staff of the Biotechnology Departments, Mr. Rosli Aslim, Mr. Azman, Mrs. Renuga a/p Panjamurti, Aluyah Marzuki, Latifah Husein and all of my post-graduates colleagues; Jameah Hamed, Noraini Abdul rahman, Ong Ming Hooi, Phang Lai Yee, Manisya Zauri, Madihah Salleh, Anisah Hassan, Normala Halimoon, Abdul Rahman, Zainal Baharum, Suhaila, Azlian, Ang, Lisa Ong, Azlina, Masdiaty, Anizam and Julia for their co-operations and help during my work.

Special thanks to my family for all of the sacrifices, time, money and encouragement's from them. The success of this project has depended on the motivation and understanding of my fellow graduates not to mention on the bounty of love showered by my family and from my dearest one. The stamina of my progress has been positively grown from day to day and it is Allah in the end of which my doa's are directed in hearts of purity so that the souls mentioned shall be granted with blessings of kindness.

Wassalam



I certify that an Examination Committee met on 6th June 2003 to conduct the final examination of Norrizan binti Abdul Wahab on her Master of Science thesis entitled "Utilisation of Selected Food Waste for the Production of Organic Acids and Bioplastics" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

ZAITON HASSAN, Ph.D.

Lecturer
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

MOHD ALI HASSAN, Ph.D


Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

MOHAMED ISMAIL ABDUL KARIM, Ph.D

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

FAKHRUL'L-RAZI AHMADUN, Ph.D

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D,
Professor/Deputy Dean,
School of Graduate Studies
Universiti Putra Malaysia

Date : 04 DEC 2003

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows :

MOHD ALI HASSAN, Ph.D

Associate Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Chairman)

MOHAMED ISMAIL ABDUL KARIM, Ph.D

Professor
Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

FAKHRUL'L-RAZI AHMADUN, Ph.D

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)




AINI IDERIS, Ph.D.
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 8 JAN 2004

DECLARATION

I hereby declare that thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other at UPM or other institutions.



(NORRIZAN ABDUL WAHAB)

Date: 3/12/03

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvii
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxi

CHAPTER

I	INTRODUCTION	1
II	LITERATURE REVIEW	4
	Household Solid Wastes.....	4
	Characteristic of Food Waste.....	5
	Utilization of Food Waste.....	8
	Biological Conversion Process.....	8
	Enzymes.....	9
	α -amylase.....	9
	Protease.....	9
	Lipase.....	10
	Cellulose.....	10
	Natural Food Fermentation.....	12
	Lactic Acid Bacteria (LAB).....	14
	Anaerobic Digestion Process.....	17
	Biochemistry.....	18
	Microbiology.....	22
	Acidogenesis.....	23
	Optimum Environmental Conditions.....	29
	pH.....	29
	Temperature.....	30
	Wastewater Analysis.....	31
	Biological Oxygen Demand (BOD).....	31
	Chemical Oxygen Demand (COD).....	32
	Culture conditions.....	33
	Batch culture.....	33
	Fed-batch culture.....	34
	Ion exchange Resin.....	35
	Converting Waste to Beneficial Materials.....	37
	The Problem with Plastics.....	38
	Polyhydroxyalkanoates (PHAs).....	39
	Structure of PHA.....	41



	Occurrence of PHA.....	45
	Environmental Conditions Affecting PHA Formation.....	48
	<i>Ralstonia Eutropha</i>	51
	Carbon Substrate.....	54
	Production Strategies.....	55
	Detection and Analytical Methods.....	57
	Extraction of Bioplastics.....	59
	Application and Commercialisation.....	60
	Economic Aspect.....	61
	Prospects.....	62
III	GENERAL MATERIALS AND METHODS.....	64
	Chemical reagents.....	64
	Food Waste.....	65
	Microorganism and Maintenance.....	66
	Preparation of Inoculum.....	66
	Preparation of Medium.....	67
	Experimental Plan.....	68
	Anaerobic Treatment Bioreactor.....	71
	Bioreactor Set-up for PHA Production.....	72
	Analytical Methods.....	74
	Quantitative Analysis.....	74
	Organic Acids Determination.....	74
	Dry Cell Weight Measurement.....	74
	Evaporation.....	75
	PHA Analyses.....	75
	Ammonium Concentration.....	76
	Chemical Oxygen Demand (COD).....	76
	Biological Oxygen Demand (BOD).....	77
	Total Solid and Ash.....	77
	Total Kjeldahl Nitrogen.....	78
	Total Sugar.....	79
	Fat.....	79
	Carbohydrate.....	80
	Fiber.....	80
	Qualitative Analysis.....	81
	Nile Blue A Staining.....	81
IV	PRODUCTION OF ORGANIC ACIDS FROM SELECTED FOOD WASTES.....	83
	Introduction.....	83
	Materials and Methods.....	84
	Food Wastes.....	84
	Substrate Preparation.....	85
	Characterization of Food Waste.....	85
	Culture Conditions.....	86
	Experimental Procedure.....	86

	Samples Analyses.....	87
	Result and Discussion.....	88
	Raw Material Analyses.....	88
	Characteristic of Food Waste.....	91
	Batch Treatment.....	95
	Effect of Substrate.....	95
	Effect of Temperature.....	99
	Effect of pH.....	104
	Conclusion.....	111
V	MICROBIOLOGICAL STUDY AND ENZYME ACTIVITY ON THE FOOD WASTE FERMENTATION.....	113
	Introduction.....	113
	Materials and Methods.....	115
	Fermentation of Food Wastes.....	115
	Viable Cell Count.....	115
	Identification of the Isolated Bacteria.....	117
	Catalase Test.....	118
	Gram-Test (Pre-screening for Gram-stain).....	118
	Carbohydrate Fermentation Profile.....	119
	Chemical Analyses.....	120
	Protease Activity (Estimation of Proteolytic Activity).....	121
	Lipase Activity.....	122
	Total Cellulose Activity.....	123
	Assay of Amylase Activity.....	124
	Result and Discussion.....	124
	Microflora Analyses.....	124
	Enzyme Activities Involved in Food Wastes Fermentation.....	134
	Total Sugars and Total acids.....	136
	Conclusion.....	138
VI	CONVERSION OF FOOD WASTES INTO BIOPLASTIC (PHB) BY <i>RALSTONIA EUTROPHA</i>.....	140
	Introduction.....	140
	Materials and Methods.....	141
	Microorganism and Culture Conditions.....	141
	Organic Acids from Food Wastes.....	142
	Culture Conditions.....	143
	Experimental Plan.....	143
	Analyses.....	145
	Polymer Extraction.....	145
	Results and Discussions.....	146
	Structure of <i>Ralstonia Eutropha</i>	146
	Nile Blue Staining.....	148
	Batch Culture Profile.....	149
	Two-stage Fed-batch Culture.....	149
	PHB Production from Synthetic Lactic Acid.....	149

	PHB Production using Concentrated Organic Acids From Food Wastes.....	151
	Effect of presence residue protein or amino acids in Media Production.....	155
	Repeated-batch Profile.....	156
	Removal of ammonium or residue protein from Organic Acids using Resins.....	158
	PHB production using ammonia-deficient organic Acids from food wastes.....	159
	General Discussions and Conclusions.....	161
VII	SUMMARY AND SUGGESTIONS FOR FUTURE WORKS.....	165
	Summary.....	165
	Suggestions for Future Work.....	168
	BIBLIOGRAPHY.....	172
	APPENDICES.....	182
	VITA.....	197

LIST OF TABLES

Table		Page
1	Characteristic of the kitchen waste.....	5
2	Comparison between food wastes taken from restaurant and households....	6
3	Nutrient composition of food waste.....	8
4	Classification of lactic acid bacteria at the generic level.....	16
5	Main genera Implicated in Anaerobic Digestion.....	22
6	Composition of cation and anion exchangers.....	36
7	Comparison between properties of polypropylene, PHB and PHB/V.....	45
8	Microbes known to accumulate PHA.....	47
9	List of limiting compounds leading to PHA formation.....	50
10	Medium composition of PHA production using <i>Ralstonia eutropha</i>	67
11	Trace element composition.....	67
12	Composition of solid defined mineral medium for Nile Blue Staining.....	82
13	Characteristic of kitchen waste (Substrate A).....	88
14	Characteristic of restaurant waste (Substrate B).....	89
15	Characteristic of restaurant waste mixed fruit waste (Substrate C).....	90
16	Composition of raw material from three kind of food waste.....	95
17	Procedure for D-form and L-form lactic acid determination.....	120
18	Comparison of PHB production using ammonia-deficient organic acids from treated food waste and synthetic medium with lactic acid.....	161



LIST OF FIGURES

Figures

1	The fermentation of glucose in homofermentative and heterofermentative lactic acid bacteria.....	15
2	Enantiomers of lactic acid.....	17
3	General scheme of the reactions occurring in anaerobic digestion.....	21
4	Reaction performed by acid producing bacteria (APB).....	25
5	End-products of various microbial fermentation.....	26
6	Pathways involved in carbohydrate fermentation by stage I fermentative bacteria.....	27
7	General structure of polyhydroxyalkanoates and some representative members.....	41
8	Electron micrograph of ultrathin sections of <i>Alcaligenes eutrophus</i> containing P(3HB) granules (white fractions). Bar, 1 μm	43
9	Metabolic pathway involved in the synthesis and degradation of poly (3-hydroxybutyrate) in <i>Alcaligenes eutrophus</i>	44
10	Biosynthesis of PHB during unbalance condition.....	51
11	General Experimental Plan.....	70
12	Bioreactor set-up for anaerobic treatment (unsteril).....	71
13	Bioreactor set-up for PHA production (sterile).....	73
14	Organic acids production from restaurant waste after 24 hours sampling.....	94
15	Organic acids produced during anaerobic treatment of kitchen garbage (substrate A) without pH control at 37°C.....	97
16	Organic acids produced during anaerobic treatment of restaurant waste (substrate B) without pH control pH at 37°C.....	97
17	Organic acids produced during anaerobic treatment of restaurant waste contain fruit waste (substrate C) without pH control at 37 °C.....	97
18	Organic acids produced during anaerobic treatment of substrate C without pH control at 30 °C.....	100

19	Organic acids produced during anaerobic treatment of substrate C without pH control at 37 °C.....	100
20	Organic acids produced during anaerobic treatment of substrate C without pH control at 45°C.....	100
21	Organic acids produced during anaerobic treatment of substrate C without pH control at 37°C.....	105
22	Organic acids produced during anaerobic treatment of substrate C initial pH 7 at 37°C.....	105
23	Organic acids produced during anaerobic treatment of substrate C controlled at pH 7 for 24 hours at 37°C.....	105
24	Organic acids produced during anaerobic treatment of substrate C controlled at pH 7 for 48 hours (2 days) at 37°C.....	107
25	Organic acids produced during anaerobic treatment of substrate C controlled at pH 7 at 37°C.....	108
26	Bacteria cell in food waste under microscope (1000x).....	125
27	Carbohydrate fermentation profile (API 50 CHL kit).....	129
28	Profile of enzyme activity during anaerobic treatment of food waste.....	133
29	Profile of total sugar related with organic acid production during anaerobic treatment of food waste.....	133
30	Scanning electron microscope (SEM) view of the species <i>Ralstonia eutropha</i> , Bar = 1µm	146
31	Transmission electron microscope (TEM) of the species <i>Ralstonia eutropha</i> is not containing polymer granules, PHA in cells.....	147
32	Transmission electron microscope (TEM) view of the accumulation of polymer granules, PHA in cells of the species <i>Ralstonia eutropha</i> (30K x 200M).....	147
33	PHA production from food waste as a carbon source by <i>Ralstonia eutropha</i>	150
34	Fed-batch fermentation by <i>Ralstonia eutropha</i> using synthetic acid as a carbon source for PHA production.....	150
35	Fed-batch fermentation by <i>Ralstonia eutropha</i> using food waste as a carbon source for PHA production.....	154

36	Studies the profile of PHB production and ammonium concentration change induced by <i>Ralstonia eutropha</i> a) Ammonia Sulfat b) Amino Acids contain.....	154
37	Repeated-batch fermentation by <i>Ralstonia eutropha</i> using organic acid from food waste as a substrate for PHA production.....	157
38	The treatment of food waste for reduces ammonium concentration using Dowex 88wx cation exchange resin in shake flask experiment.....	157
39	Fed-batch fermentation by <i>Ralstonia eutropha</i> using treatment organic acids from food waste as a carbon source for PHA production.....	160

LIST OF ABBREVIATIONS

BOD ₅	- Biochemical oxygen demand (after 5 days incubation of sample at 20°C in the dark)
C	- Concentration
C/N	- Carbon to nitrogen ratio
CO ₂	- Carbon dioxide
COD	- Chemical Oxygen demand
GC	- Gas chromatography
HPLC	- High pressure liquid chromatography
ICI	- Imperial Chemical Industries
PHAs	- Polyhydroxyalkanoates
P(3HB)	- Poly (3-hydroxybutyrate)
PHBV	- Poly (hydroxybutyrate-co-hydroxyvalerate)
VFA	- Volatile fatty acid
LAB	- Lactic acid bacteria
TCA	- Trichloroacetic acid
CHO	- Carbohydrate

CHAPTER 1

INTRODUCTION

Food waste is on the increase with bigger population and affluence. Food waste is non-toxic, rich in organic matter and highly biodegradable. Most of these wastes end up in landfill sites, whereby the carbonaceous compounds are eventually decomposed to biogas. Apart from contributing to environmental pollution, such a system destroys the organic matter, which could be utilized as useful materials for producing value added products such as organic acids. Food wastes contain carbohydrates, proteins and lipids. These compounds are essential for the growth of microorganisms to synthesize desired products. Through biological treatment, food waste can be subjected for organic acids production, particularly for lactic acid production under controlled conditions. The organic acids can then be used as substrates by microbes to produce bioplastics.

It is known that many bacteria can produce biopolymers intracellularly under certain growth conditions, such as excess carbon and limiting nitrogen. These biopolymers have properties similar to normal plastics, with the advantage that they are biodegradable upon disposal in the soil within a few months. Through research, alternatives for energy-intensive chemical processes can be explored, thus reducing our national dependence on petroleum. Developing biodegradable plastic is one way to solve the waste disposal problem. Biodegradable plastics that do not pollute the global environment has become an extremely important product (Doi, 1990).

These biodegradable plastic materials must retain the desired material properties of conventional synthetic plastic, and should be completely degraded without leaving any undesirable residues when discarded. However, such bioplastics are not widely used currently due to their high costs. Currently, PHA in the form of poly (3HB-co-3HV) or PHB/V is being produced on a fairly large scale by Imperial Chemical Industries Limited (ICI) in Britain under the trade name Biopol using *Alcaligenes eutrophus* H16. Unfortunately, the high cost of PHA has limited their use on a wide scale. ICI's Biopol is sold for more than US\$4.40/kg compared to polypropylene, which is only US\$0.60/kg. This high cost is mainly due to the carbon source and the use of large quantities of organic solvents in the recovery process for extracting the intracellular polymers (Hassan, 1997).

The high PHA production cost can possibly be lowered by using unpurified, low cost organic wastes from agriculture and food processing plants (Cho *et al.*, 1997). In this study, different compositions of food waste were chosen as substrates, which are rich in organic compounds necessary for bacterial growth. PHA or biodegradable thermoplastic polyhydroxyalkanoates can be produced from food waste in a two-stage process. During the first stage, the complex organic compounds are digested under anaerobic conditions by bacteria into volatile fatty acids (VFA) such as acetic, lactic, butyric and propionic acids. Then, in the second stage, the PHA-producing bacteria such as *Ralstonia eutropha* (previously known as *Alcaligenes eutrophus*) can utilize the VFA and polymerize the acids into PHA as carbon and energy reserves under nitrogen limited conditions.

Objectives

The objectives of this study are as follows:

- ❑ To obtain the best conditions such as substrate composition, temperature and pH in the utilization of food waste for the production of organic acids as materials for the production of PHA.
- ❑ To identify microbial and enzymatic activities affecting the production of organic acids during the fermentation of food waste.
- ❑ To produce polyhydroxyalkanoate (PHA) by *Ralstonia eutropha* using organic acids from anaerobically treated food waste.